


AP Chemistry Exam Review

Free Response Practice #2

2018 #6 (4 points)



A student sets up a galvanic cell at 298 K that has an electrode of Ag(s) immersed in a 1.0 M solution of Ag⁺(aq) and an electrode of Cr(s) immersed in a 1.0 M solution of Cr³⁺(aq), as shown in the diagram above.

- a. The student measures the voltage of the cell shown above and discovers that it is zero. Identify the missing component of the cell, and explain its importance for obtaining a nonzero voltage.

Half-Reaction	E° (V)
$\text{Ag}^+(\text{aq}) + e^- \rightarrow \text{Ag}(\text{s})$	+ 0.80
$\text{Cr}^{3+}(\text{aq}) + 3 e^- \rightarrow \text{Cr}(\text{s})$?

- b. The student adds the missing component to the cell and measures E°_{cell} to be +1.54 V. As the cell operates, Ag⁺ ions are reduced. Use this information and the information in the table above to do the following.
- Calculate the value of E° for the half-reaction $\text{Cr}^{3+} + 3 e^- \rightarrow \text{Cr}(\text{s})$.
 - Write the balanced net-ionic equation for the overall reaction that occurs as the cell operates.
 - Calculate the value of ΔG° for the overall cell reaction in J/mol_{rxn}.